

Magnetic Quasiparticles in sQGP

Explain Low Viscosity and Static Potentials

Jinfeng Liao^{a,1} and Edward Shuryak^{a,2}

^a Department of Physics & Astronomy, SUNY Stony Brook, NY 11794, USA.

¹ jiliao@ic.sunysb.edu, ² shuryak@tonic.physics.sunysb.edu

Although “dual superconductor” nature of confinement was suggested by 't Hooft and Mandelstam long ago, the importance of magnetic quasiparticles near deconfinement transition and their relation to the “perfect liquid” observed at RHIC were realized only last year [1,2]. In the RHIC region $T=1-2T_c$ both electric and magnetic quasiparticles are important degrees of freedom. They are about equally massive and abundant at $1.4T_c$ and below this T the magnetic ones get lighter, more weakly coupled, and become the dominant component.

One evidence of this phenomenon is magnetic screening mass which exceeds electric one. Another is the persistence of “flux tube”-related linear potential, seen in potential energy and entropy associated with static quarks on the lattice. In a model of electric-magnetic plasma [3,4] we explained its tension — which is few times larger than at $T = 0$ and is seen till about $1.3T_c$ — and fix the magnitude of the magnetic pressure $P_M(T)$ in this region. The condition for electric flux tube’s existence/disappearance is derived: it led to magnetic particle density $n \sim 4.4 - 6.6/fm^3$ at $1.3T_c$, consistent with several lattice results.

We found that a plasma made of electric and magnetic quasiparticles leads to mutual trapping, similar to “magnetic bottle effect”, and explains low viscosity and small diffusion observed at RHIC. We used Molecular Dynamics (MD) for simulation of a strongly coupled plasma consisting of totally ~ 1000 charges of both types with variable density ratio of electric/magnetic components. The measured shear viscosity and diffusion constant are the lowest in the maximally mixed plasma (i.e. 50%-50%) which takes the most advantage of trapping effect, and their values after suitable mapping to sQGP system’s parameter regions fall very close to those suggested by RHIC experiments. The importance of this effect for understanding such short mean free path as indicated by RHIC is further elucidated in [5].

If this mechanism is the only one, and since it is operative only not too far from T_c , then viscosity/diffusion will grow away from “perfect liquid” regime above $2T_c$ and LHC would show significantly different picture than RHIC.

References

- [1] E. Shuryak, “Strongly Coupled Quark-Gluon Plasma: The Status Report”, *arXiv: hep-ph/0608177*.
- [2] J. Liao and E. Shuryak, “Strongly Coupled Plasma with Electric and Magnetic Charges”, *Phys. Rev. C*, **75**, (2007) 054907.
- [3] J. Liao and E. Shuryak, “Electric Flux Tube in Magnetic Plasma”, *arXiv:0706.4465*.
- [4] J. Liao and E. Shuryak, “String Tension at Finite Temperature and Magnetic Pressure on Electric Object”, *in preparation*.
- [5] J. Liao and E. Shuryak, “Lorentz Trapping Makes Liquid with Low Viscosity and Small Diffusion”, *in preparation*.